

§ 558.105

Sections Affected in the Finding Aids section of this volume.

§ 558.105 [Reserved]

§ 558.115 Carbadox.

(a) *Approvals*. Type A medicated articles: 2.2 percent (10 grams per pound) to 000069 in § 510.600(c) of this chapter.

(b) *Related tolerances*. See § 556.100 of this chapter.

(c) *Special considerations*. Do not use in Type B or Type C medicated feeds containing bentonite.

(d) *Conditions of use*. It is used for swine as follows:

(1) *Amount per ton*. 10–25 grams (0.0011–0.00275 percent).

(i) *Indications for use*. For increase in rate of weight gain and improvement of feed efficiency.

(ii) *Limitations*. Do not feed to swine weighing more than 75 pounds body weight; do not feed to swine within 10 weeks of slaughter; do not use in complete feeds containing less than 15 percent crude protein.

(2) *Amount per ton*. 50 grams (0.0055 percent).

(i) *Indications for use*. For control of swine dysentery (vibrionic dysentery, bloody scours, or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); increased rate of weight gain and improved feed efficiency.

(ii) *Limitations*. Do not feed to swine weighing more than 75 pounds body weight; do not feed to swine within 10 weeks of slaughter; do not use in complete feeds containing less than 15 percent crude protein.

(3) *Amount per ton*. Carbadox 50 grams (0.0055 percent) plus pyrantel tartrate, 96 grams (0.0106 percent).

(i) *Indications for use*. For control of swine dysentery (vibrionic dysentery, bloody scours, or hemorrhagic dysentery); control of bacterial swine enteritis (salmonellosis or necrotic enteritis caused by *Salmonella choleraesuis*); aid in the prevention of migration and establishment of large roundworm (*Ascaris suum*) infections; aid in the prevention of establishment of nodular worm (*Oesophagostomum*) infections.

(ii) *Limitations*. Do not feed to swine over 75 pounds; do not feed within 10 weeks of slaughter; consult a veteri-

21 CFR Ch. I (4–1–98 Edition)

narian before feeding to severely debilitated animals; feed continuously as sole ration. Do not use in complete feeds containing less than 15 percent crude protein.

[40 FR 13959, Mar. 27, 1975, as amended at 40 FR 45164, Oct. 1, 1975; 40 FR 57798, Dec. 12, 1975; 42 FR 761, Jan. 4, 1977; 51 FR 7396, Mar. 3, 1986]

§ 558.120 Carbarsone (not U.S.P.).

(a) *Approvals*. Type A medicated articles: (1) 37.5 percent to 046573 in § 510.600(c) of this chapter.

(2) 25 percent carbarsone and 5 grams per pound bacitracin (as bacitracin methylene disalicylate) to 046573 in § 510.600(c) of this chapter.

(b) *Related tolerances*. See § 556.60 of this chapter.

(c) [Reserved]

(d) *Conditions of use*. (1) It is used for turkeys as follows:

(i) *Grams per ton*. 227 to 340.5 (0.025 to 0.0375 percent).

(a) *Indications for use*. As an aid in the prevention of blackhead.

(b) *Limitations*. Feed continuously beginning 2 weeks before blackhead is expected and continue as long as prevention is needed; withdraw 5 days before slaughter; as sole source of organic arsenic.

(ii) *Grams per ton*. 227 to 340.5 (0.025 to 0.0375 percent) carbarsone plus 10 grams per ton bacitracin from bacitracin methylene disalicylate.

(a) *Indications for use*. As an aid in the prevention of blackhead; for increased rate of weight gain.

(b) *Limitations*. Feed continuously beginning 2 weeks before blackhead is expected and continue as long as prevention is needed; withdraw 5 days before slaughter; as sole source of organic arsenic.

(iii) *Grams per ton*. 227 to 340.5 (0.025 to 0.0375 percent) carbarsone plus 4 to 45 grams per ton bacitracin from bacitracin zinc.

(a) *Indications for use*. As an aid in the prevention of blackhead, increased rate of weight gain, and improved feed efficiency.

(b) *Limitations*. Feed continuously as sole ration. Withdraw 5 days before slaughter. As sole source of organic arsenic; as bacitracin zinc provided by